



Richmond Pediatrics
 357 NW Richmond Beach Road
 Shoreline, WA 98177
 (206) 546-2421

ACETAMINOPHEN (Tylenol)

		Suspension	Children's Chews	Jr. Strength Chews
weight (lbs)	weight (kg)	5 ml = 160 mg	1 tab = 80mg	1 tab = 160 mg
< 6 lbs	< 2.7 kg	Consult Provider		
6-11 lbs	2.7-5.3 kg	1.25 ml		
12-17 lbs	5.4-8 kg	2.5 ml	1 tab	
18-23 lbs	8.1-10.9 kg	3.75 ml	1 1/2 tabs	
24-35 lbs	11- 16.3 kg	5 ml	2 tabs	1 tab
36-47 lbs	16.4-21.7 kg	7.5 ml	3 tabs	1 1/2 tabs
48-59 lbs	21.8- 27.2 kg	10 ml	4 tabs	2 tabs
60-71 lbs	27.3- 32.6 kg	12.5 ml	5 tabs	2 1/2 tabs
72-95 lbs	32.7-43.2 kg	15 ml	6 tabs	3 tabs
96+ lbs	43+ kg	20 ml	8 tabs	4 tabs

Tylenol (acetaminophen) can be given every 4-6 hours, not more than 5 doses in 24 hours.

*Infant and children's suspension have the same concentration but come with different measuring tools.

IBUPROFEN (Motrin or Advil)

weight (lbs)	weight (kg)	Infant Drops*	Children's Suspension*	Jr. Strength Chews
do not give to age < 6 mo		1.25 ml = 50mg	5 ml = 100 mg	1 tab = 100mg
< 11 lbs	< 5.4 kg	Consult Provider		
12-17 lbs	5.4-8 kg	1.25 ml	2.5 ml	
18-23 lbs	8.1-10.9 kg	1.875 ml	3.75 ml	
24-36 lbs	11- 16.3 kg	2.5 ml	5 ml	1 tab
36-47 lbs	16.4-21.7 kg		7.5 ml	1 1/2 tabs
48-59 lbs	21.8- 27.2 kg		10 ml	2 tabs
60-71 lbs	27.3- 32.6 kg		12.5 ml	2 1/2 tabs
72-95 lbs	32.7-43.2 kg		15 ml	3 tabs
96+ lbs	43+ kg		20 ml	4 tabs

Ibuprofen (Motrin, Advil) may be given every 6-8 hours, no more than 4 doses in 24 hours.

*Infant suspension and children's liquid have DIFFERENT strengths.